

Scotland's Rural College

Xylanase and xylo- oligosaccharide prebiotic improve the growth performance and concentration of potentially prebiotic oligosaccharides in the ileum of broiler chickens

Craig, AC; Khattak, FM; Hastie, P; Bedford, Michael R; Olukosi, OA

Published in:
British Poultry Science

DOI:
[10.1080/00071668.2019.1673318](https://doi.org/10.1080/00071668.2019.1673318)

Print publication: 01/02/2020

Document Version
Peer reviewed version

[Link to publication](#)

Citation for pulished version (APA):

Craig, AC., Khattak, FM., Hastie, P., Bedford, M. R., & Olukosi, OA. (2020). Xylanase and xylo- oligosaccharide prebiotic improve the growth performance and concentration of potentially prebiotic oligosaccharides in the ileum of broiler chickens. *British Poultry Science*, 61(1), 70-78. <https://doi.org/10.1080/00071668.2019.1673318>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Xylanase and xylo- oligosaccharide prebiotic improve the growth performance and concentration of potentially prebiotic oligosaccharides in the ileum of broiler chickens

Journal:	<i>British Poultry Science</i>
Manuscript ID	CBPS-2019-183.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	05-Aug-2019
Complete List of Authors:	<p>Craig , Allison ; Scotland's Rural College, Monogastric Science Research Centre; University of Glasgow, McCall Building, School of Veterinary Medicine</p> <p>Khattak, Farina ; Scotland's Rural College, Monogastric Science Research Centre</p> <p>Hastie, Peter ; University of Glasgow, McCall Building, School of Veterinary Medicine</p> <p>Bedford, Michael; ABVista, Woodstock Centre, Marlborough Business Park</p> <p>Olukosi, Oluyinka; Scotland's Rural College; University of Georgia, Department of Poultry Science</p>
Keywords:	Broilers, Xylanase, Nutrient digestibility, Xylose oligosaccharides, short chain fatty acids, Growth

SCHOLARONE™
Manuscripts

Xylanase and xylo-oligosaccharide prebiotic improve the growth performance and concentration of potentially prebiotic oligosaccharides in the ileum of broiler chickens.

A. D. Craig^{1,2}, F. Khattak¹, P. Hastie², M.R. Bedford³, and O. A. Olukosi^{1,4*}.

¹ Monogastric Science Research Centre, SRUC, Edinburgh, EH9 3JG

²McCall Building, School of Veterinary Medicine, University of Glasgow, G61 1QH

³A B Vista, Woodstock Centre, Marlborough Business Park, Marlborough, SN8 4AN

⁴ Department of Poultry Science, University of Georgia, Athens, GA 30602, USA

*Corresponding author

Oluyinka A. Olukosi, Department of Poultry Science, University of Georgia, Athens, GA 30602 Email: oaolukosi@uga.edu Tel: +1 706 542 1359

ORCID- Miss A.D Craig – 0000-0002-1640-8612

ORCID- Dr O.A Olukosi- 0000-0002-9137-6745

ABSTRACT

1. The objective of this study was to investigate the effect of supplementing broiler diets with xylanase or xylo- oligosaccharide (XOS) on growth performance, the concentration of non-starch polysaccharide (NSP) hydrolysis products in the ileum and concentration of short chain fatty acids (SCFA) in the caeca of broiler chickens.
2. In total, 500 male Ross 308 broilers were used in this 29-day (d) study. The treatments were organised into a 2×2 plus 1 factorial arrangement consisting of two additives (xylanase or XOS) at two levels (low or high) plus a control treatment with no additives. This gave five treatments with 100 bird in each treatment group. The diets were slightly deficient in protein by 20 g/kg and energy by 1 MJ/kg.
3. On d 14 and 28, two birds per pen were euthanised, the caeca content collected and analysed for short chain fatty acid (SCFA) concentration. On d 29, six birds per pen were euthanised and ileal digesta were collected and analysed for the concentration of NSP fractions.
4. On d 14, caecal acetic acid, iso-butyric acid, iso-valeric acid, n-valeric acid and total SCFA concentrations were significantly greater ($P \leq 0.05$) when diets were supplemented with XOS compared with xylanase.
5. Ileal concentration of arabinose, galactose and glucuronic acid (GlucA2) were significantly greater ($P \leq 0.05$) in the insoluble NSP fraction when diets were supplemented with a high level of xylanase, compared with the control treatment. Ileal concentration of fructose was significantly greater ($P \leq 0.05$) in the water soluble NSP when a high level of xylanase or low level of XOS were included in the diet compared with the control.

6. It was concluded that xylanase and XOS had similar effects on NSP concentration and SCFA in the caeca, although there was little effect on performance. This observation demonstrated further benefits of xylanase supplementation in wheat-based broiler diets beyond digesta viscosity reduction and the release of extra nutrients.

Keywords- xylanase, XOS, broilers, performance, nutrient digestibility, SCFA

50 Introduction

51 Depression in growth performance caused by an increase in digesta viscosity is a
52 common occurrence in broiler diets containing a large amount of non-starch
53 polysaccharides (NSP; Jia *et al.*, 2009). In order to overcome this, carbohydrase
54 enzymes are often added to broiler diets to improve nutrient utilisation and increase
55 productivity. Carbohydrases hydrolyse NSP, breaking it down into smaller
56 oligosaccharides. This results in a decrease in digesta viscosity and the release of
57 encapsulated nutrients (Knudsen, 2014). In addition to these benefits, it has been
58 suggested that the small oligosaccharides produced during NSP hydrolysis could
59 have prebiotic properties (Courtin *et al.*, 2008).

60 One way of investigating the production of small oligosaccharides is to measure the
61 concentration of NSP hydrolysis products in the ileum of broilers, such as arabinose
62 or xylose concentrations. A prebiotic is a small molecule which is fermented by
63 beneficial bacteria, encouraging their growth, while discouraging the growth of
64 pathogenic bacteria. Xylo-oligosaccharides (XOS) are associated with improvements
65 in poultry performance (Al-Sultan *et al.*, 2016) by modulating the gastrointestinal
66 immune system, microbial populations (Jung *et al.*, 2008; Yang *et al.*, 2008) and
67 increasing short chain fatty acid (SCFA) production, however, evidence for this has
68 been inconsistent (Arsi *et al.*, 2015; Gao *et al.*, 2007).

69 To investigate the production of potentially prebiotic oligosaccharides during
70 xylanase activity, monosaccharides were measured in the ileal digesta and compared
71 to monosaccharides in digesta from birds supplemented with purified XOS. The
72 objective of the trial was to investigate similarities in profiles of monosaccharides in

73 digesta from birds receiving xylanase and those receiving XOS to illustrate that NSP
74 hydrolysis products may have prebiotic-like effects similar to that of purified XOS.
75 NSP hydrolysis products are thought to be fermented by beneficial bacteria such as
76 *Bifidobacter* and *Lactobacilli spp.*, producing SCFA (Lee *et al.*, 2017). An increase
77 in the concentration of SCFAs is often associated with an increase in the population
78 of beneficial bacteria and a decrease in pathogenic bacteria (Engberg *et al.*, 2004). In
79 addition to this, SCFAs have been shown to influence growth performance in
80 broilers. Butyrate, in particular, is regarded as an available energy source, increasing
81 the energy available to the host for growth (Ravangard *et al.*, 2017). Supplementing
82 broiler diets with xylanase or XOS has been shown to affect the production of SCFA
83 in the caeca of broiler chickens (Engberg *et al.*, 2004; Lee *et al.*, 2017). This could
84 suggest that both ingredients have a similar mode of action. As such, the effect of
85 xylanase or XOS on SCFA concentration in the caeca of broilers was investigated in
86 the current study.

87 The objective of this experiment was to investigate the effect of supplementing
88 wheat-based diets, which were deficient in energy and protein, with xylanase or XOS
89 on growth performance, the concentration of NSP hydrolysis products in the ileum
90 and the concentration of SCFA's in the caeca of broiler chickens.

92 **Materials and methods**

93 *Animals and management*

94 All the procedures in the experiment were approved by the SRUC Animal
95 Experiment Committee.

Five hundred male Ross 308, one-d-old broilers were allocated to one of five treatments organised as a randomised complete block design. The birds were housed 10 in a pen, with ten pen replicates per treatment and provided with feed and water on an *ab libitum* basis throughout the experiment (0 to 29 d). The treatments followed a 2×2 plus 1 factorial arrangement with two different additives (xylanase or XOS) at two inclusion levels (high and low) plus the control. The low level of xylanase inclusion was 16, 000 XU/kg and the high was 32,000 XU/kg. The low level of XOS inclusion was 0.25 g/kg and the high level was 0.1 g/kg. The lower level of each additive was based on the standard recommendation by the manufacturer while the higher level of each additive was chosen following a literature search (Wang *et al.*, 2005; Zhenping *et al.*, 2013; De Measschalck *et al.*, 2015). The control diet was deficient in energy by 1 MJ/kg and protein reduced by 3%, to 20g/kg CP however; all other nutrient requirements were met to ensure that any effects were induced by energy or protein deficiency or additive supplementation alone. The diets were formulated to be deficient in energy and protein to allow any improvements in growth or nutrient digestibility to become apparent, as previous work has indicated that xylanase supplementation may be more beneficial in nutrient-deficient diets (Francesch and Geraert, 2009). The experiment was split into five treatments; 1) control, 2) control plus xylanase 16,000 XU/kg, 3) control plus xylanase 32,000 XU/kg, 4) control plus purified XOS 0.25 g/kg and 5) control plus purified XOS 1.0 g/kg. All of the experimental diets were provided in mash form and birds had *ad libitum* access to feed and water throughout the feeding trial. The xylanase used contained 160 000 U of endo- 1,4 β xylanase activity per gram. One unit of xylanase activity was defined as the amount of enzyme required to liberate 1 μmol of reducing sugars from xylan using a standardised test (Enzyme

121 Services (ESC), Hengoed, Ystrad Mynack, UK). The xylanase (Econase XT) was
122 supplied by AB Vista, Marlborough, UK. The XOS used in this study was purchased
123 from Shandong Lifelong Bio-technology Co., China (XOS 35A). The ingredient and
124 chemical composition of the control diet is shown in Table 1. Wheat bran was
125 included in the diet during this study to increase the amount of NSP in the diet and
126 maximise the potential for prebiotic oligosaccharide generation during NSP
127 hydrolysis by xylanase.

128

129 Table 1 here

130

131 ***Growth performance***

132 Feed and birds were weighed on d 0, 14 and 28. The data from feed and bird weights
133 were used to calculate body weight gain (BWG), feed intake (FI) and feed
134 conversion ratio (FCR).

135 ***Sample collection***

136 On d 14 and 28, two birds per pen were euthanised by cervical dislocation and used
137 to collect caeca content for SCFA analysis. Following euthanasia, the caeca were
138 removed and the content was gently squeezed into a collection tube.

139 On d 29, the remaining six birds per pen were euthanised by cervical dislocation and
140 ileal content was collected for nutrient digestibility and NSP analysis. Once located,
141 the distal half of the ileum was removed and the contents were flushed with water
142 into a collection pot. The ileal digesta from all six birds per pen was pooled and was

143 collected on day 29, as opposed to day 28, due to practical reasons relating to
144 volume of samples to be collected.

145 ***Short chain fatty acid analyses***

146 Once the contents of the caeca were removed they were stored at -20°C and later
147 analysed for SCFA concentration as described by Khattak *et al.* (2018) using gas
148 chromatography.

149 ***NSP fraction analyses***

150 The ileal digesta samples collected from six birds per pen on d 29 were dried in a
151 Unitherm force draft drying oven for 48 hours at 80°C. The samples were analysed
152 for water extractable (WE) carbohydrate components and water unextractable (WU)
153 NSP using HPLC, following the method of Englyst *et al.* (1994). The pre-caecal
154 NSP concentration was calculated using the equation displayed in the calculations
155 section.

156 ***Chemical analysis***

157 The ileal digesta samples collected on d 29 were dried prior to conducting titanium
158 and DM analysis according to the method of Short *et al.* (1996). Dry matter (DM)
159 was determined using standard methods from AOAC, whereby 1g of the sample was
160 dried in a uniform forced drying oven (Unitherm, Russel-Lindsay Engineering Ltd,
161 Birmingham, England, UK) at 95°C for 24 hours. Nitrogen determination was
162 carried out by the combustion method (Method 934.01; AOAC). Gross energy was
163 determined using an isoperibol bomb calorimeter system using benzoic acid as an
164 internal standard (Model 6200, Parr Instruments, Moline, Illinois, USA). Ileal
165 digestibility was calculated using the index method described by Olukosi *et al.*

(2007). The activity of xylanase in the diet was measured using a commercial test kit (Enzyme Services (ESC), Hengoed, Ystrad Mynack, UK). One BXU was defined as the amount of xylanase enzyme required to liberate 1 nmol of reducing sugars per minute from xylan at pH 5.3.

Calculations

Pre-caecal NSP concentration (g/100g DM intake) was calculated using the equation below:

$$NSP\ Concentration = NSP\ conc.\ in\ digesta \times \left(\frac{Ti\ in\ diet}{Ti\ in\ digesta} \right)$$

NSP Conc. In digesta (%)

Statistics

Statistical analysis was conducted using the ANOVA function of Genstat (16th Edition). Data were analysed following the 2 × 2 plus 1 factorial arrangement with body weight as the blocking factor. When an interaction between additive type and inclusion level (excluding the no additive treatment) was significant, the means for growth performance and nutrient digestibility were separated using Tukey's test. The significant additive type × inclusion level interactions for ileal NSP concentration were detected using specific contrasts details of which are in Tables 5 and 6. Significance was set at P≤0.05 and trends at P<0.1.

Results

Growth performance of broilers on days 14 and 28

187 The enzyme analysis results showed that diets 1, 4 and 5 contained xylanase activity
188 below detection threshold. Diet 2 contained 16,600 BXU/kg and diet 3 contained
189 36,900 BXU/ kg of xylanase activity.

190 On d 14 there was no additive type×inclusion level interaction for BWG, FI or FCR
191 both on d 14 and 28 (Table 2). However, there were main effects of additive type and
192 inclusion level on d 14. Body weight gain and feed intake were greater ($P<0.05$)
193 following xylanase supplementation compared with XOS supplementation. Feed
194 conversion ratio was lower ($P<0.05$) following additive supplementation at high
195 inclusion level compared with low inclusion level. There was no effect of additive
196 supplementation compared with the unsupplemented control.

197 On d 28, there were no significant ($P<0.05$) main or interaction effects of additive
198 type or inclusion level for BWG, FI or FCR. However, feed intake and FCR values
199 were lower ($P<0.001$) for broilers receiving xylanase or XOS compared with the
200 control.

201

202 Table 2 here

203

204 ***Nutrient digestibility in broilers aged 29 days and fed diets supplemented with***
205 ***xylanase or XOS***

206 Nutrient digestibility in broilers aged 29 d and fed diets supplemented with xylanase
207 or XOS is shown in Table 3. There was a significant additive type×inclusion level
208 interaction for nitrogen (N) intake. Birds receiving diets containing the low level of
209 xylanase had significantly ($P<0.01$) lower N intake than those receiving the high

210 level of xylanase. There was no effect of additive type or inclusion level on DM or
211 N digestibility, however, there was trend for greater ileal digestible energy (IDE) in
212 birds fed diets supplemented with xylanase compared to those with XOS. The DM,
213 IDE and N digestibilities were lower ($P<0.05$) in birds fed diets containing xylanase
214 or XOS compared to the control. The N and gross energy intake were lower ($P<0.05$)
215 in birds fed diets containing xylanase or XOS compared to the control diets.

216

217 Table 3 here

218

219 ***Short chain fatty acid concentration in the caeca of broiler chickens on days 14***
220 ***and 28***

221 Caecal SCFA concentration in response to xylanase or XOS supplementation on d
222 14 is shown in Table 4. There was no significant inclusion level or additive
223 type×inclusion level interaction for SCFA concentration on d 14, however there was
224 a main effect of ingredient type. The concentration of acetic acid, propanioc acid,
225 iso-butyric acid, iso- valeric acid and total SCFA were greater ($P<0.05$) following
226 XOS compared with xylanase supplementation. There was no effect of ingredient
227 inclusion compared to the control treatment on acetic acid, n-butyric acid or n-
228 valeric acid concentration in broilers on d 14. However, there was a trend ($P=0.066$)
229 for xylanase to decrease propionic acid concentration and XOS to increase propionic
230 acid concentration.

231 Caecal SFCFA concentration in response to xylanase or XOS supplementation on d
232 28 is shown in Table 4. There was no additive type×inclusion level interaction for

233 SCFA concentration on d 28. There was no main effect of ingredient type or
234 inclusion level on caecal SCFA concentration on day 28. The concentration of acetic
235 acid, n-butyric acid and total SCFA were greater ($P<0.05$) following ingredient
236 supplementation compared to the control. The concentration of acetic acid,
237 propanioc acid and total SCFA was greater ($P<0.05$) in birds aged 28 days compared
238 to birds aged 14 days.

239

240 Table 4 here

241

242 ***NSP fraction content of ileal digesta from broiler chickens aged 29 days***

243 The concentration of WU NSP fractions in response to xylanase or XOS
244 supplementation is shown in Table 5. There was an ingredient type \times inclusion level
245 interaction for arabinose and galactose. WU arabinose and galactose concentration
246 were greater when diets were supplemented with the high level of xylanase
247 compared to the low level. When the control treatment was compared to the other
248 treatments individually, arabinose and galactose concentration were greater ($P<0.05$)
249 following xylanase supplementation at a high level, compared to the control. There
250 were main effects of ingredient type and inclusion levels. Rhamnose concentration
251 was greater ($P<0.001$) following XOS compared to xylanase supplementation.
252 Rhamnose and fructose concentration were greater ($P<0.05$) following
253 supplementation at a high level compared to the low level. The concentration of
254 rhamnose, fructose, arabinose and galactose were greater ($P<0.05$) following
255 supplementation compared with the control.

256

257 Table 5 here

258

259 The concentration of WE NSP fractions in response to xylanase or XOS
260 supplementation is shown in Table 6. There was a significant ingredient
261 type×inclusion level interaction for fructose concentration. The WE fructose
262 concentration was greater when the high level of xylanase or the low level of XOS
263 were included in the diet, compared to the low level of xylanase or high level of
264 XOS. When the control treatment was compared to the other diets individually, WE
265 fructose concentration was greater ($P<0.05$) following supplementation of 32,000
266 XU/kg xylanase or 0.025% XOS compared to the control. There were no main
267 effects of ingredient type or inclusion level for WE NSP concentration. There was a
268 tendency ($P=0.061$) for greater xylose concentration in diets supplemented with XOS
269 compared to xylanase. There was a tendency for greater galactose concentration in
270 diets supplemented with the high compared to the low inclusion level. Galactose and
271 total WE NSP concentration were greater ($P<0.05$) following supplementation
272 compared with the control.

273

274 Table 6 here

275

276 **Discussion**

277 Xylanase is used routinely in broiler diets to improve growth performance, however,
278 there is evidence to suggest that potentially prebiotic oligosaccharides are generated
279 during NSP hydrolysis. The generation of *in-situ* prebiotics could provide additional
280 benefit to the use of xylanases in broiler diets over and above the reduction in digesta
281 viscosity.

282 ***Growth Performance***

283 The birds in the current study performed below breed standards. This could be due to
284 a multitude of reasons, including diet form and composition. It has been well
285 established in the literature that low protein (23 -20% CP) or low energy (3000-
286 2640 Kcal/kg) diets reduce the growth performance of broilers (Govil *et al.*, 2017;
287 Kamran *et al.*, 2008; Williams *et al.*, 2014). The current study is in agreement with
288 this, as the birds receiving the control feed, which was lower in energy by 8% and
289 protein by 13%, ate significantly more than those birds receiving supplementation in
290 their diet. This was expected to an extent, as it has been suggested that birds do not
291 eat to satisfy hunger *per se* but they consume enough feed to satisfy their energy or
292 protein requirements (Karam *et al.*, 2008). In the current study, the birds receiving
293 diets low in energy and protein were able to increase their feed intake, allowing them
294 to maintain their growth. When xylanase or XOS was added into the diet, feed intake
295 reduced, as expected, resulting in a reduction in FCR and an increase in efficiency.

296 Xylanase supplementation improved BWG, FI and FCR compared to that of XOS on
297 d 14. This was expected as the mechanism of action of xylanase is well established
298 in the literature. Xylanase cleaves the arabinoxylan (AX) backbone of NSP releasing
299 the trapped nutrients (Meng *et al.*, 2005) and reduces digesta viscosity (Lentle,
300 2005). This impacts on growth performance in two ways. Firstly, the release of

trapped nutrients increases the amount available for absorption in the small intestine (Meng *et al.*, 2005). Secondly, reducing digesta viscosity allows sufficient mixing of the digesta, enabling more nutrients to be absorbed (Lentle, 2005). There is evidence to suggest a third mechanism, namely generation of *in-situ* prebiotic oligosaccharides. During the hydrolysis of NSP, smaller oligosaccharides such as XOS are generated which have been shown to have prebiotic-like effects (Zhang *et al.*, 2014).

Prebiotics can improve growth performance of broilers (Al-Sultan *et al.*, 2016; Abdel- Hafeez *et al.*, 2017). In the current study, on d 28, XOS reduced feed intake and improved FCR but had no effect on body weight gain, which was similar to the effect of xylanase supplementation. It is thought that prebiotics improve growth performance by increasing nutrient absorption due to modulating gut microflora and increasing gut integrity (Al-Sultan *et al.*, 2016). Beneficial gut bacteria, such as *Bifidobacterium* and *Lactobacilli spp.*, ferment prebiotics, such as XOS, FOS and GOS, which encourages growth whilst discouraging the colonisation of pathogenic bacteria (Xu *et al.*, 2003; Courtin *et al.*, 2008; Yousaf *et al.*, 2016). This may relate to the SCFA results, discussed below.

Nutrient digestibility

Ingredients such as xylanase have been shown to improve nutrient digestibility (Kiarie *et al.*, 2014). The expectation that adding carbohydrases can improve nutrient digestibility is logical, as the enzyme should decrease digesta viscosity and allow increased absorption of nutrients (Mathlouthi *et al.*, 2002). In the current study, DM and N digestibility were reduced when either xylanase or XOS were added to the diet, however, IDE was significantly lower when XOS was included in the diet. The

325 diets used during this study were deficient in energy and protein, so the aim of
326 adding such ingredients was to improve nutrient absorption, as this is what has been
327 described in the literature (Cowieson *et al.*, 2017). The addition of xylanase or XOS
328 reduced digestible energy and N intake compared to the control treatment, which
329 may help explain why nutrient digestibility was decreased following
330 supplementation.

331 The negative effect of XOS supplementation on IDE was unexpected. The literature
332 often reports no effect of prebiotics on ileal nutrient digestibility (Kirkpinar *et*
333 *al.*, 2004; Mountzouris *et al.*, 2010) however improvements in total tract retention
334 have been reported (Mountzouris *et al.*, 2010). This is achieved when populations of
335 beneficial microflora are encouraged, which increases nutrient digestion and
336 adsorption (De Measschalck *et al.*, 2015) of SCFAs produced during fibre
337 fermentation. Decreased nutrient digestibility following prebiotic supplementation
338 has been reported previously in pigs. The authors suggested that the reduction in
339 nutrient digestibility was due to the introduction of indigestible fibre into the diet
340 (Smiricky- Tjardes *et al.*, 2003). This is an unlikely explanation for the reduction in
341 nutrient digestibility reported in the current study, however, the inclusion level of XOS
342 was much lower than that used by Smiricky- Tjardes *et al.*, (2003).

343 There was an effect of supplementation on nutrient digestibility however there was
344 little effect on growth performance especially on d 28, meaning that the
345 improvements in nutrient digestibility were not translated into growth performance,
346 which has been reported previously (Yang *et al.*, 2008; Gonzalez- Ortiz *et al.*, 2016).
347 This could indicate that 'point in time' measurements, such as nutrient digestibility,
348 are poor tools in predicting the long-term effects of a diet on performance parameters
349 such as body weight gain or FCR.

350 ***Short chain fatty acid concentration***

351 Bird age had a significant effect on the concentration of SCFAs in the caecum of
352 broiler chickens. On d 28, there was a greater concentration of acetic, propionic, iso-
353 butyric and iso-valeric acid compared to SCFA on d 14. This is in agreement with
354 Lee *et al.* (2017). The authors showed that the concentration of SCFAs increased as
355 the bird aged. The reason for this was likely to be the development of the intestinal
356 microflora. Gong *et al.*, (2008) demonstrated that young birds (14 d of age) had a
357 less well-developed microflora than those aged 42 d. Not only was the microflora in
358 14 d old birds less well developed, it was more likely to be influenced by changes in
359 the diet or environment. In the current study, it was possible that SCFA
360 concentration in the caeca were lower on d 14, because the microflora were less well
361 developed and not able to ferment carbohydrate sources in order to produce SCFA.
362 As the bird aged, the microflora matured and established itself, enabling the
363 microbes to more readily ferment available carbohydrates, increasing the production
364 of SCFA, which is in agreement with Lee *et al.*, (2017).

365 The differences in SCFA concentration may have been related to differences in
366 growth recorded in the current study, especially on d 14. SCFAs can influence
367 growth performance in different ways. Firstly, as previously mentioned, SCFA can
368 be used as an energy source for colonic cells, which increases the nutrients available
369 to the host for growth. Secondly, xylanase enzymes randomly cleave NSP, reducing
370 digesta viscosity and increasing nutrient absorption resulting in a decrease in the
371 amount of nutrients available to the micro-organisms the caeca for fermentation (Lee
372 *et al.*, 2017).

373 One of the most notable detrimental nutrients fermented by colonic bacteria is
374 protein. Protein fermentation by colonic bacteria has been associated with the
375 production of toxic compounds, such as ammonia, which can inhibit growth and
376 even cause disease (Apajalahil and Vinenola, 2016). To combat this, it has been
377 recommended that xylanases should be used to increase nutrient utilisation and to
378 create fermentable carbohydrates (Apajalahil and Vinenola, 2016). The current data
379 agreed with this, as growth performance and SCFA production increased following
380 the xylanase supplementation, however there was no effect on nitrogen digestibility.

381 ***Pre- caecal NSP fraction concentration***

382 NSP concentration is a way to measure its hydrolysis by giving an indication of the
383 resulting sugars in the liquid and solid phase of the ileal digesta. In a previous study,
384 WU NSP concentration decreased while WE NSP concentration increased following
385 xylanase supplementation (Olukosi *et al.*, 2015). A reduction in NSP concentration
386 indicates greater hydrolysis, which is beneficial when xylanase is used to reduce
387 digesta viscosity. In the current study, however, the aim was to increase the
388 concentration of potentially prebiotic oligosaccharides and investigate their effect on
389 growth performance. NSP from wheat contains large amount of arabinoxylan (AX).
390 WE NSPs are the main cause of increased digesta viscosity in broilers, which is
391 counteracted by using carbohydrases such as xylanase (Choct, 2015). The
392 concentration of WE NSP in the ileum of broilers during the current study increased
393 when either xylanase or XOS were supplemented in the diets. This is in agreement
394 with previous published data, as it has been reported that the disappearance of NSP
395 from the digestive tract of broilers was significantly affected by enzyme addition
396 (Cowieson *et al.*, 2016; Cozannet *et al.*, 2017). From the data above, fructose and

397 galactose concentration (which likely represented fructo- oligosaccharide (FOS and
398 galacto- oligosaccharide (GOS) or galactans) was significantly increased due to
399 xylanase addition. In addition to this, xylose concentration tended to increase XOS
400 addition.

401 This tendency for xylose to increase following XOS supplementation could be as a
402 result of undigested XOS bypassing digestion, but stoichiometry suggested this
403 could not have been the only source of xylan in the WE fraction as 0.1 g/kg was
404 added to the diet, and concentration increased by 3 g/ kg DM. An increase in
405 galactose (representing galactans and pectins) concentration in the ileum of broilers
406 following carbohydrase supplementation has been reported previously (Kocher *et al.*,
407 2002). This is of interest, because FOS and GOS have been associated with prebiotic
408 effects (Kaplan and Hutkins, 2000; Boehm *et al.*, 2005; Courtin *et al.*, 2008).

409 WU NSP are generally not fermented, unlike WE NSP; however, they have been
410 associated with gut development (Choct, 2015). In the current study, the
411 concentration of arabinose, fructose and galactose were increased in response to
412 xylanase or XOS supplementation in the WU NSP fraction. This could suggest that
413 xylanase may have beneficial effects on gut development which is in agreement with
414 other studies (Jimenez –Monero *et al.*, 2009; Liu *et al.*, 2017). This is certainly true
415 for the gizzard, where increases in dietary fibre have been associated with heavier
416 gizzards, resulting in increased retention time of digesta and smaller particle size
417 which, in turn, increases nutrient digestibility (Jimenez- Monero *et al.*, 2009). The
418 increase in WU NSP concentration could improve the development of the
419 gastrointestinal tract, contributing to improvements in growth performance.

420 The current trial data demonstrated similarities between the effects of xylanase and
421 XOS supplementation on growth performance, pre-caecal NSP concentration and
422 caecal SCFA concentration, indicating similar modes of action. Consequently, it can
423 be suggested that improvements in growth performance were partly driven by the
424 production of in-situ prebiotics following xylanase supplementation. This study
425 suggested that supplementing broiler diets with low levels of xylanase or high levels
426 of XOS had a positive effect on the concentration of WU NSP fractions in the ileum,
427 however, there was a limited effect of high levels of xylanase or XOS on
428 performance. As such, more research into increasing the inclusion level of xylanase
429 or XOS is required.

430

431 **Acknowledgements**

432 The authors acknowledge the staff and students at the Monogastric Science Research
433 Centre, Ayr for all of their help and support.

434 Allison Craig is a recipient of i-CASE PhD studentship from Biotechnology and
435 Biological Sciences Research Council (BBSRC). SRUC receives funding from the
436 Scottish government (RESAS).

437 **Disclosure statement**

438 M.R Bedford is an employee of AB Vista, the manufacturer of the enzymes used in
439 this study.

440

441 **References**

- 442 ABDEL- HAFEEZ, H.M, E.S.E. SALEH, S. S.TAWFEEK, I.M.I YOUSSEF and
443 A.S.A
- 444 ABDEL- DAIM. 2017. “Effects of Probiotic, Prebiotic and Symbiotic With
445 and Without Feed Restriction on Performance, Haematological Indices and
446 Carcass Characteristics of Broiler Chickens”. *Asian- Australasian Journal of*
447 *Animal Science* 30: 672-682.
- 448 AL-SULTAN, S., S.M. ABDEL- RAHEEM , W.R. EL- GHAREEB and M.H.A
449 MOHAMED. 2016. “Comparative Effects of Using Prebiotic, Probiotic,
450 Symbiotic and Acidifier on Growth Performance, Intestinal Microbiology
451 and Histomorphology of Broiler Chicks”. *Japanese Journal of Veterinary*
452 *Research* 64 (supplement 2): S187-195.
- 453 APAJALAHIL, J., and K. VINENOLA. 2016. “Interaction Between Chicken
454 Intestinal
455 Microflora and Protein Digestion”. *Animal Feed Science and*
456 *Technology* 221: 323-330.
- 457 ARSI, K, A.M DONOGUE, A. WOO-MING, P.J. BLORE and D.J DONOGUE.
458 2015.
459 “The Efficacy of Selected Probiotic and Prebiotic Combinations in Reducing
460 Campylobacter Colonisation in Broiler Chickens”. *Journal of Applied*
461 *Poultry Research* 24: 327-334.
- 462 AOAC 2006. “Official Methods for Analysis, 18th ed”. *Association of Official*
463 *Analytical Chemists, Washington, DC*.
- 464 BOEHM, G, B. STAHL, J. JELINEK, J. KNOL, V. MINIELLO and G.E. MORO.
465 2005.

- 466 “Prebiotic Carbohydrates in Human Milk and Formulas”. *Aceta Paediatrica*
 467 94 (suppl 449): 18-21.
- 468 CHOCT, M. 2015. “Feed Non-Starch Polysaccharides for Monogastric Animals:
 469 Classification and Function”. *Animal Production Science*. 27: 855-861.
- 470 COURTIN, C.M, W.F. BROEKAERT, K. SWENEN, O. LESCROART, O.
 471 ONAGBESAN, J. BUYSE, E. DECUYPERE, T. VAN DEWIELE, M.
 472 MARZORATI, W. VERSTRATE, G. HUYGHEBAERT and J.A.
 473 DELCOUR. 2008. “Dietary Inclusion of Wheat Bran
 474 Arabinooxylooligosaccharides Induces Beneficial Nutritional Effects in
 475 Chickens”. *Cereal Chemistry*. 85: 607-613.
- 476 COWIESON, A.J, H. LU, K.M. AJUWON, I. KNAP and O. ADEOLA. 2017.
 477 “Interactive Effects of Dietary Protein Source and Exogenous Protease and
 478 Jejunal Health of Broiler Chickens”. *Animal Production Science*. 57: 252-
 479 261.
- 480 COWIESON, A.J, W. SCHLIFKA, I. KNAPP, F.F. ROOS, R. SCHOOP and J.W.
 481 WILSON. 2016. “Meta- Analysis of Effect of a Mono-Component Xylanase
 482 on the Nutritional Value of Wheat Supplemented with Exogenous Phytase
 483 for Broiler Chickens”. *Animal Production Science*. 56: 2014- 2022.
- 484 COZANNET, P, M.T KIDD, R.M. NETO and P.A. GERAERT. 2017. “Next-
 485 Generation Non-Starch- Polysaccharide- Degrading, Multi Carbohydrase
 486 Complex Rich in Xylanase and Arabinofuranosidase to Enhance Feed
 487 Digestibility”. *Poultry Science*. 96: 2743-2750.
- 488 De MEASSCHALCK, C, V. EECKHAUT, L. MAERTENS, L. DE LAUGE, L.
 489 MARCHAL, C. NEZER, S. De BAERE, S. CROUBLES, G. DAUBE, J.
 490 DEWULF, F. HAESBROUCK, R. DUCATELLE, B. TAMINAU and F.

- 491 VAN IMMERSEEL. 2015. "Effects of Xylo-oligosaccharides on Broiler
492 Chicken Performance and Microbiota". *Applied and Environmental*
493 *Microbiology* 81: 5880-5888.
- 494 ENGBERG, R.M, M.S. HEDENANN, S. STEINFELDT AND B.B JENSEN. 2004.
495 "Influence of Whole wheat and Xylanase on Broiler Performance and
496 Microbial Composition and Activity in the Digestive Tract". *Poultry Science*
497 82: 925-938.
- 498 ENGLYST, H. N., M.E. QUIGLEY AND G. J. HUDSON. 1994. "Determination of
499 Dietary Fibre as Non-Starch Polysaccharides with Gas-Liquid
500 Chromatographic, High-Performance Liquid Chromatographic or
501 Spectrophotometric Measurement of Constituent Sugars". *Analysts* 119:
502 1497-1509.
- 503 FRANCESCH. M AND P.A GERAERT. 2009. " Enzyme Complex Containing
504 NSPase and Phytase Improves Growth Performance and Bone Mineralisation
505 of Broilers Fed Reduced Nutrient Corn-Soybean- Based Diets". *Poultry*
506 *Science* 88; 1915-1924.
- 507 GAO, F., Y. JIANG, G.H. ZHOU, AND Z.K. HAN. 2007. "The Effects of Xylanase
508 Supplementation on Growth, Digestion, Circulating Hormone and Metabolite
509 Levels, Immunity and Gut Microflora in Cockerels Fed on Wheat Based
510 Diets". *British Poultry Science* 48: 480-488.
- 511 GONG J, H. YU, T. LIU, J.J GILL, J.R. CHAMBERS, R. WHEATCROFT AND
512 P.M.
513 SABOUR. 2008. "Effects of Zinc Bacitracin, Bird Age and Access to Range
514 on Bacterial Microbiota in the Ileum and Caeca of Broiler Chickens". *Journal*
515 *of Applied Microbiology* 104 :1372-1382.

- 516 GONZALEZ- ORTIZ, G, O. OLUKOSI AND M.R BEDFORD. 2016. "Evaluation
517 of
518 the Effect of Different Wheats and Xylanase Supplementation on
519 Performance, Nutrient and Energy Utilisation in Broiler Chicks". *Animal*
520 *Nutrition* 2; 173-179.
- 521 GOVIL, K., S. NAYAK, R.P.S BAGHEL, A.K. PATIL, C.D.MALAPURE and D.
522 THAKUR. 2017. "Performance of Broiler Chicken Fed Multi-Carbohydrases
523 Supplemented Low Energy Diet". *Veterinary World* 10: 727-731.
- 524 JIA, W., B.A. SLOMINSKI, H.L. BRUCE, G. BLANK, G. CROW and O. JONES.
525 2009. "Effects of Diet Type and Enzyme Addition on Growth Performance
526 and Gut Health of Broiler Chickens During Sub Clinical *Clostridium*
527 *perfringens* Challenge". *Poultry Science* 88: 132-140.
- 528 JIMENEZ- MONERO, E, J.M. GINZALEZ-ALVARADO, A. DECOCA- SINOVA,
529 R.
530 LAZARO and G.G. MATEOS. 2009. "Effects of Source of Fibre on the
531 Development and pH of the Gastrointestinal Tract of Broilers". *Animal Feed*
532 *Science and Technology* 154: 93-101.
- 533 JUNG, S.J, R. HOUDE, B. BAURHOO, X. ZHAO and B.H. LEE. 2008. "Effects of
534 Galacto- oligosaccharides and a Bifidobacteria Lactic-based Probiotic Strain
535 on the Growth Performance and Faecal Microflora of Broiler Chickens".
536 *Poultry Science* 87: 1694-1699.
- 537 KAMRAN, Z., M. SARWAR, M. NISA, M.A. NADEEM, S.MAHMOOD,
538 M.E.BABAR,
539 and S. AHMED. 2008. "Effect of Low Protein Diets Having Constant
540 Energy-to- Protein Ratio on Performance and Carcass Characteristics of

- 541 Broiler Chickens From One to Thirty- Five Days of Age”. *Poultry Science*
542 87: 468-474.
- 543 KAPLAN, H and R.W. HUTKINS. 2000. “Fermentation of Fructooligosaccharides
544 by
545 Lactic Acid Bacteria and Bifidobacteria”. *Applied and Environmental*
546 *Microbiology* 66: 2682-2684.
- 547 KHATTAK, F, V. PASCHALIS, M. GREEN, J.G.M HOUDJIK, P. SOULTANAS
548 AND
549 J. MAHDAVI. 2018. “TYPLEX® Chelate, a Novel Feed Additive, Inhibits
550 *Campylobacter jejuni* Biofilm Formation and Cecal Colonization in Broiler
551 Chickens”. *Poultry Science* 97: 1319-1399.
- 552 KIARIE, E, L.F. ROMERO and V. RAVINDRAN. 2014. “Growth Performance,
553 Nutrient Utilisation and Digesta Characteristics in Broiler Chickens Fed
554 Corn or Wheat Diets Without or With Supplemental Xylanase”. *Poultry*
555 *Science* 93: 1186-1196.
- 556 KIRKPINAR, K, Z. ACIKGOZ, M. BOZKURT and V. AYHAN. 2004. “Effects of
557 Inclusion of Poultry By-product Meal and Enzyme- Prebiotic
558 Supplementation in Grower Diets on Performance and Feed Digestibility of
559 Broilers”. *British Poultry Science* 45: 273-279.
- 560 KNUDSEN, K.E.B. 2014. “Fibre and Non-Starch Polysaccharide Content and
561 Variation in Common Crops used in Broiler Diets”. *Poultry Science* 93:
562 2380-2393.
- 563 KOCHER, A, M. CHCOT, M.D. PORTER and J. BROZ. 2002. “Effect of Feed
564 Enzymes on the Nutritive Value of Soybean Meal Fed to Broilers”. *British*
565 *Poultry Science* 43: 54-63.

- LEE, S.A, J. APAJALAHTI, K. VIENOLA, G. GONZALEZ-ORTIZ, C.M.G.A.
FONTES and M.R. BEDFORD. 2017. "Age and Dietary Xylanase
Supplementation Affects Ileal Sugar Residues and Short Chain Fatty Acid
Concentration in the Ileum and Caecum of Broiler Chickens". *Animal Feed
Science and Technology* 234: 29-42.
- LENTLE, R.G. 2005. "The Macro-biophysics of Digestion; Implications for the
Poultry Industry". *Australian Poultry Science Symposium* 17: 163-170.
- LIU, W.C and I.H. KIM. 2017. "Effects of Dietary Xylanase Supplementation on
Performance and Functional Digestive Parameters in Broilers fed Wheat-
based Diets". *Poultry Science* 96: 566-573.
- MATHLOUTHI, N, S. MALLET, L. SAULINER, B. QUEMENER and M. Larbier.
2002. "Effects of Xylanase and β -glucanase Addition on Performance,
Nutrient Digestibility and Physico Chemical Conditions in the Small
Intestine Contents fed a Wheat and Barley Based Diet". *Animal Research* 51:
395- 406.
- MENG, X. and B.A. SLOMINSKI. 2005. "Nutritive Values of Corn, Soybean Meal,
Canola Meal and Peas for Broiler Chickens as Affected by a
Multicarbohydrase Preparation of Cell Wall Degrading Enzymes". *Poultry
Science* 84: 1242-1251.
- MOUNTZOURIS, K. C., P. TSITRSIKOS, I. PALAMIDI, A. ARVANITI, M.
MOHNL,
G. SCHATZMAYR and K. FEGEROS. 2010. "Effects of Probiotic
Inclusion Levels in Broiler Nutrition on Growth Performance, Nutrient
Digestibility, Plasma Immunoglobulins, and Caecal Microflora
Composition". *Poultry Science* 89: 58-67.

- 591 OLUKOSI, O. A., A.J. COWIESON and O. ADEOLA. 2007. "Age-related
592 Influence
593 of a Cocktail of Xylanase, Amylase, and Protease or Phytase Individually or
594 in Combination in Broilers". *Poultry Science* 86: 77–86.
- 595 OLUKOSI, O.A, L.A. BEESON, K. ENGLYST and L.F. ROMERO. 2015. "Effects
596 of
597 Proteases Without or With Carbohydrases on Nutrient Digestibility and
598 Disappearance of Non- Starch Polysaccharides in Broiler Chickens". *Poultry
599 Science* 94: 2662-2669.
- 600 RAVANGARD, A.H, M. HOUSHMAND, M. KHAJVI AND R. NAGHIHI. 2017.
601 "Performance and cecal bacteria counts of broilers fed low protein diets with
602 and without a combination of probiotic and prebiotic". *Brazilian Journal of
603 Poultry Science*. Special Issue- Nutrition: 75-82
- 604 SHORT, F.J, P. GORTON, J. WISEMAN and K.N. BOORMAN. 1996.
605 "Determination of Titanium Oxide Added as an Inert Marker in Chicken
606 Digestibility Studies". *Animal Feed Science Technology* 59: 215-221.
- 607 SMIRICKY- TJARDES, M.R, C.M. GRIESSHOP, E.A. FLICKINGER, L.L
608 BAUER
609 and G.C. Fahey Jr. 2003. "Dietary Galactooligosaccharides Affect Ileal and
610 Total Tract Nutrient Digestibility, Ileal and Faecal Bacterial Concentrations
611 and Ileal Fermentative Characteristics of Growing Pigs". *Journal of Animal
612 Science* 81: 2535-2545.
- 613 WANG, Z.R, S.Y QIAO, W.Q LU and D.F LI. 2005. "Effects of enzyme

- 614 supplementation on performance, nutrient digestibility, gastrointestinal
615 morphology and volatile fatty acid profiles in the hindgut of broilers fed
616 wheat based diets”. *Poultry Science* 84: 875-881.
- 617 WILLIAMS, M.P, J.T. KLEIN, C.L. WYATT, T.W. YORK and J.T. Lee. 2014.
618 “Evaluation of Xylanase in Low Energy Broiler Diets”. *Journal of Applied*
619 *Poultry Research* 23: 188-195.
- 620 XU, Z.R., G.H. HU, M.S. XIA, X.A. ZHAN and M.Q. Wang. 2003. “Effects of
621 Dietary
622 Fructooligosaccharide on Digestive Enzyme Activities, Intestinal Microflora
623 and Morphology of Male Broilers”. *Poultry Science* 82: 1030-1036.
- 624 YANG, Y, P.A. IJI, L.L MIKKELSEN and M. Choct. 2008. “Effects Dietary
625 Mannanligosaccharide on Growth performance and Gut Development of
626 Broilers Given Different Cereal- Based Diets”. *Journal of Animal Physiology*
627 *and Animal Nutrition* 92: 650-659.
- 628 YOUSAF, M.S, A. IJAZ, K. ASHRQAF, M.A. RASHID, A. HAFEEZ, H. ZANEB,
629 E.
630 DAR, R. NASEER, I. RABBANI, J. ZENTEK and H. Reham. 2016.
631 “Comparative Effects of Different Dietary Concentrations of β -galacto-
632 oligosaccharides on Growth Performance, Feed Efficiency and Organ
633 Development in Broiler Chickens”. *The Journal of Animal and Plant Science*
634 26: 1603-1608.
- 635 ZHANG, L, J. XU, L. LEI, Y. JIANG, F. GAO and G.H. ZhOU. 2014. “Effects of
636 Xylanase Supplementation on Growth Performance, Nutrient Digestibility
637 and Non-Starch Polysaccharide Degradation in Difference Sections of the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

638 Gastro-intestinal Tract of Broilers”. *Asian-Australasian Journal of Animal*
639 *Science* 27: 855-861.

640 ZHENPING, S, L WENTING, Y RUIKUI, L JIA, L HONGHONG, S WEI, W
641 ZHONGMIE, L JINGPAN, S ZHE and Q YULING. 2013. “Effect of straw
642 derived oligosaccharide on broiler growth performance, endocrine
643 metabolism and immune response”. *Canadian Journal of Veterinary*
644 *Research* 77: 105-109.

645

Table 1. The calculated and analysed nutrient content of wheat-based diets supplemented with enzymes or prebiotic oligosaccharide.

Ingredient (g/kg)	Control
Wheat	556.5
Wheat Bran	192.5
Soybean meal	152.5
Soya oil	25.0
Limestone	12.5
Dicalcium Phos, 18%P	14.0
Sodium Bicarbonate	6.0
Lysine HCl	5.0
Methionine	2.0
Threonine	2.0
Valine	2.0
Vitamin & Mineral premix	5.0
TiO ₂ Marker	25.0
Total	1000.0
Calculated content	
ME (MJ/kg)	11.41
Crude Protein (g/kg)	200.0
Calcium (g/kg)	10.0
Phosphorus (g/kg)	7.5
nPP (g/kg)	4.9
Na (g/kg)	2.1
Cl (g/kg)	3.5
Analysed nutrient content	
DM (g/kg)	877.1
AME (MJ/kg)	16.19
Na (g/kg)	1.9
Cl (g/kg)	3.1
Calcium (g/kg)	9.6
Phosphorus (g/kg)	6.3
N(g/kg)	31.3

Notes; TiO₂- titanium dioxide; Na- sodium; Cl- chloride; DM- dry matter; AME- apparent metabolizable energy; N- nitrogen

Table 2. The growth performance of broilers fed diets deficient in energy and protein and supplemented with xylanase or XOS.

Additive	Inclusion Level	Day 14			Day 28		
		BWG (g/bird)	FI (g/bird)	FCR	BWG (g/bird)	FI (g/Bird)	FCR
No Additive		388.1	596.2	1.54	1313.3	2953.9	2.26
Xylanase	Low	381.1	594.1	1.56	1294.5	2707.1	2.10
	High	394.3	579.2	1.47	1320.0	2546.5	1.93
XOS	Low	356.7	561.5	1.59	1261.6	2536.5	2.03
	High	375.1	561.7	1.51	1306.4	2604.7	2.00
	SEM	10.449	12.08	0.038	29.202	86.161	0.078
P-value of control vs additive ¹		0.390	0.111	0.752	0.592	<0.001	0.009
Means for main effect of additive type (AT)							
Xylanase		387.7 ^b	586.7 ^b	1.52 ^a	1307.3	2626.8	2.02
XOS		365.9 ^a	561.6 ^a	1.55 ^b	1284.0	2570.6	2.02
	SEM	10.449	12.08	0.038	29.202	86.161	0.078
Means for main effect of inclusion level (IL)							
	Low	368.9	577.8	1.57	1278.0	2621.8	2.07
	High	384.7	570.4	1.49	1313.2	2570.6	1.97
	SEM	10.449	12.08	0.038	29.202	86.161	0.078
P values							
Additive Type (AT)		0.044	0.045	0.449	0.430	0.518	0.978
Inclusion Level (IL)		0.139	0.543	0.034	0.236	0.595	0.197
AT × IL		0.806	0.535	0.944	0.744	0.193	0.385

Notes; XOS- xylo-oligosaccharide; BWG- body weight gain; FI- feed intake; FCR; feed conversion ratio; ¹P- values for control vs. additive types - the mean for the control group was compared to all other treatments containing an additive irrespective of type or level ; ^{ab} different superscripts with the same column indicate means that are significantly (P<0.05) different

Table 3. Coefficients of energy and nitrogen digestibility of broilers aged 29 days and fed diets supplemented with xylanase or XOS

Additive	Inclusion Level	DM	IDE (MJ/kg)	N	Energy Intake (units/bird)	N Intake (units/bird)
No Additive		0.713	13.77	0.858	38.44	85.37
Xylanase	Low	0.652	13.28	0.837	33.23	81.18 ^b
	High	0.606	11.99	0.817	29.62	69.15 ^a
XOS	Low	0.630	12.22	0.812	29.54	70.23 ^{ab}
	High	0.624	12.23	0.821	30.26	79.64 ^{ab}
	SEM	0.0189	0.372	0.00913	1.499	1.987
P-value of control vs additive ¹		<0.001	0.003	0.001	<0.001	0.002
Means for main effect of additive type (AT)						
Xylanase		0.629	12.63	0.827	31.39	75.71
XOS		0.627	12.23	0.816	29.94	74.39
	SEM	0.0189	0.263	0.00646	1.06	1.987
Means for main effect of inclusion level (IL)						
	Low	0.641	12.75	0.824	31.43	75.16
	High	0.615	12.11	0.819	29.90	74.94
	SEM	0.0189	0.263	0.00646	1.06	1.405
P values						
Additive Type (AT)		0.170	0.095	0.550	0.317	0.937
Inclusion Level (IL)		0.909	0.284	0.245	0.343	0.643
AT × IL		0.302	0.090	0.115	0.158	<0.001

Notes; XOS- xylo-oligosaccharide; DM- dry matter; IDE- ileal digestible energy; N- nitrogen; ¹P- values for control vs. additive types - the mean for the control group was compared to all other treatments containing an additive irrespective of type or level; ^{abc} different superscripts with the same column indicate means that are significantly (P<0.05) different

Table 4. The effect feeding wheat-based diets supplemented with xylanase or XOS on SCFA concentrations (mg/kg) in the caeca on days 14 and 28

Additive	Inclusion Level	Day 14					Day 28				
		Acetic Acid	Propionic Acid	n- butyric Acid	n- Valeric Acid	Total SCFA	Acetic Acid	Propionic Acid	n- butyric Acid	n- Valeric Acid	Total SCFA
No Additive		4749	75.70	1260.7	45.3	6242	4790	193	1046	69.1	6170
Xylanase	Low	4260	56.60	1117.8	23.1	5532	5284	198	1347	78.2	6984
	High	4036	61.0	1250.0	28.7	6438	5541	172	1563	80.8	7435
XOS	Low	4949	112.0	1348.6	40.6	6528	5374	166	1345	80.1	7056
	High	5244	136.4	1505.9	55.0	7018	5177	222	1194	82.9	6756
Pooled SEM		108.8	35.54	143.13	9.87	437.2	234.9	28.9	114.8	6.92	310.6
P-value of control vs. Additive ¹		0.737	0.066	0.874	0.596	0.818	0.042	0.922	0.019	0.149	0.015
Means for main effect of additive types (AT)											
Xylanase		4148 ^a	56.80 ^a	1183.9	32.4 ^a	5485 ^a	5412	185	1455	79.5	7210
XOS		5096 ^b	124.2 ^b	1427.3	47.8 ^b	6773 ^b	5275	194	1269	81.5	6906
Pooled SEM		236.5	47.66	172.07	10.89	309.1	116.1	20.5	81.1	4.89	219.6
Means for main effect of inclusion level (IL)											
Low		4605	82.30	1233.2	38.4	6030	5412	185	1455	79.5	7210
High		4640	98.70	1378.0	41.9	6228	5275	194	1269	81.5	6906
Pooled SEM		236.5	11.60	102.35	2.47	309.1	116.1	20.5	81.1	4.89	219.6
P- values for main effects and interactions											
Additive Type (AT)		0.007	<0.001	0.128	0.026	0.006	0.563	0.760	0.114	0.774	0.335
Inclusion Level (IL)		0.917	0.265	0.335	0.759	0.654	0.899	0.612	0.778	0.699	0.808
AT × IL		0.442	0.546	0.789	0.234	0.509	0.341	0.168	0.118	0.989	0.234

Notes; XOS- xylo-oligosaccharide; ¹P- values for control vs. additive types - the mean for the control group was compared to all other treatments containing an additive irrespective of type or level; ^{ab} different superscripts with the same column indicate means that are significantly (P<0.05) different

Table 5. The effect of feeding wheat-based diets supplemented with xylanase or XOS on the concentration (g/100g) of WU NSP fractions in the ileum of broilers

Additive	Inclusion Level	Rhamnose	Fructose	Arabinose	Xylose	Galactose	TOTAL (g/100g)
No Additive		0.0156	0.0184	1.647 ^a	2.54	0.583 ^a	8.74
Xylanase	Low	0.0129	0.0170	1.721 ^a	2.65	0.572 ^a	9.01
	High	0.0222	0.0374	2.431 ^b	3.55	0.851 ^b	12.13
XOS	Low	0.0261	0.0256	2.002 ^{ab}	3.08	0.686 ^{ab}	10.30
	High	0.0434	0.0450	1.950 ^{ab}	2.90	0.688 ^{ab}	10.35
	Pooled SEM	0.004	0.005	0.155	0.292	0.047	0.840
P-value of control vs additive types ¹		0.022	0.044	0.042	0.142	0.039	0.086
Means for main effect of additive type (AT)							
Xylanase		0.0176 ^a	0.0272	2.076	3.10	0.711	10.57
XOS		0.0347 ^b	0.0353	1.976	2.99	0.687	10.32
	Pooled SEM	0.003	0.004	0.110	0.206	0.033	0.594
Means for main effect of inclusion level							
	Low	0.0195 ^a	0.0213 ^a	1.861 ^a	2.87	0.629 ^a	9.65
	High	0.0328 ^b	0.0412 ^b	2.190 ^b	3.22	0.770 ^b	11.24
	Pooled SEM	0.003	0.004	0.110	0.206	0.033	0.594
P- values for main effects and interactions (IL)							
Additive Type (AT)		<0.001	0.144	0.525	0.717	0.608	0.776
Inclusion Level (IL)		0.002	0.001	0.047	0.235	0.007	0.074
AT × IL		0.302	0.937	0.024	0.083	0.008	0.084
P- Values for Contrasts							
Control vs. Xylanase, low level				0.739		0.864	
Control vs. Xylanase, high level				0.002		<0.001	
Control vs. XOS, low level				0.122		0.139	

Control vs. XOS high level	0.184	0.130
----------------------------	-------	-------

Notes; XOS- xylo-oligosaccharide; GlucA2- glucuronic acid; ¹P-value for control vs. additive types- the mean for the control group was compared to all other treatments containing an additive irrespective of type or level; ^{ab} different superscripts within the same column indicate means that are significantly different

For Peer Review Only

Table 6. The effect of feeding wheat based diets supplemented with xylanase or XOS on the concentration (g/100g) of WE NSP fractions in the ileum of broilers

Additive	Inclusion Level	Rhamnose	Fructose	Arabinose	Xylose	Galactose	TOTAL (g/100g)
No Additive		0.0390	0.0462	0.605	0.948	0.398	2.74
Xylanase	Low	0.0422	0.0479 ^a	0.624	0.898	0.463	3.04
	High	0.0619	0.0692 ^b	0.855	1.143	0.607	3.84
XOS	Low	0.0484	0.0682 ^b	0.773	1.284	0.507	3.74
	High	0.0520	0.0570 ^a	0.810	1.216	0.556	3.63
	Pooled SEM	0.010	0.007	0.08	0.115	0.052	0.351
P-value of control vs additive types ¹		0.278	0.083	0.081	0.162	0.030	0.051
Means for main effect of additive types (AT)							
Xylanase		0.0520	0.0586	0.739	1.02	0.535	3.44
XOS		0.0502	0.0626	0.791	1.25	0.532	3.68
	Pooled SEM	0.007	0.005	0.055	0.081	0.037	0.248
Means for main effect of inclusion level (IL)							
	Low	0.0453	0.0580	0.699	1.091	0.485	3.39
	High	0.0569	0.0631	0.832	1.179	0.581	3.73
	Pooled SEM	0.007	0.005	0.055	0.081	0.037	0.248
P- values for main effects and interactions							
Additive Type (AT)		0.850	0.573	0.511	0.061	0.949	0.495
Inclusion Level (IL)		0.245	0.479	0.103	0.453	0.078	0.337
AT × IL		0.471	0.032	0.227	0.190	0.369	0.261
P- values for contrasts							
Control vs. Xylanase, low level			0.866				
Control vs. Xylanase, high level			0.032				
Control vs. XOS, low level			0.039				
Control vs. XOS high level			0.291				

Notes; XOS- xylo-oligosaccharide; GlucA2- glucuronic acid; ¹P-value for control vs. additive types- the mean for the control group was compared to all other treatments containing an additive irrespective of type or level; ^{ab} different superscripts within the same column indicate means that are significantly (P<0.05) different

For Peer Review Only